

DISCOVERY OF A NEW PHOTOMETRIC SUB-CLASS OF FAINT AND FAST CLASSICAL NOVAE

M. M. KASLIWAL¹, S. B. CENKO², S. R. KULKARNI¹, E. O. OFEK¹, R. QUIMBY¹, A. RAU³*Draft version March 10, 2010*

ABSTRACT

We present photometric and spectroscopic follow-up of a sample of extragalactic novae discovered by the Palomar 60-inch telescope during a search for “Fast Transients In Nearest Galaxies” (P60-FasTING). Designed as a fast cadence (1-day) and deep ($g < 21$ mag) survey, P60-FasTING was particularly sensitive to short-lived and faint optical transients. The P60-FasTING nova sample includes 10 novae in M31, 6 in M81, 3 in M82, 1 in NGC 2403 and 1 in NGC 891. This significantly expands the known sample of extragalactic novae beyond the Local Group, including the first discoveries in a starburst environment. Surprisingly, our photometry shows that this sample is quite inconsistent with the canonical Maximum Magnitude Rate of Decline (MMRD) relation for classical novae. Furthermore, the spectra of the P60-FasTING sample are indistinguishable from classical novae. We suggest that we have uncovered a sub-class of faint and fast classical novae in a new phase space in luminosity-timescale of optical transients. Thus, novae span two orders of magnitude in both luminosity and time. Perhaps, the MMRD, which is characterized only by the white dwarf mass, was an over-simplification. Nova physics appears to be characterized by quite a rich four-dimensional parameter space in white dwarf mass, temperature, composition and accretion rate.

Subject headings: stars: novae, cataclysmic variables, galaxies: starburst, galaxies: individual (M81, M82, M31, NGC 2403, NGC 891), techniques: spectroscopic, techniques: photometric, surveys

1. INTRODUCTION

Since the discovery of classical novae, astronomers have pursued their use as standard candles to determine distances (see Hubble 1929). Zwicky (1936) first noticed some regularity in nova light curves and termed this the “life-luminosity” relation. Arp (1956) undertook a comprehensive search for novae in M31, discovering thirty novae in 290 nights, and found a clear relation — luminous novae evolve faster than less luminous novae. The modern name for this observation is the maximum-magnitude rate-of-decline relation (MMRD relation).

The MMRD relation has attracted considerable theoretical attention (e.g. Livio 1992). The basic idea is that the relation is entirely due to the mass of the accreting white dwarf. The more massive the white dwarf, the higher the surface gravity, the higher the pressure at the base of envelope and stronger the thermonuclear runaway (and hence, higher the peak luminosity). Also, the more massive the white dwarf, the smaller the envelope mass to attain the critical pressure for thermonuclear runaway (TNR) and hence, faster the decline.

In more recent times, della Valle & Livio (1995) used a sample of novae in M31 and LMC to propose an arc-tangent relation between the peak luminosity and rate of decline. Downes & Duerbeck (2000) used a sample of Galactic novae to propose a linear relation between the same two parameters. Darnley et al. (2006) used a score of novae in M31 from POINT AGAPE survey and claimed their observations were consistent with the

della Valle & Livio (1995) formulation of the MMRD.

In comparison to supernovae, classical novae are not very luminous. Hence, searches have traditionally focussed only on the Milky Way and its nearest neighbors (Andromeda and the Large Magellanic Cloud). Hornoch et al. (2008) looked into archival data and found 49 nova candidates⁴ in M81 in the past 20 years — unfortunately, these candidates neither have light curves nor spectra. Ferrarese et al. (2003) undertook a search for novae using 24 orbits of the Hubble Space Telescope and found nine nova candidates in M49. Even with their sparsely sampled light curves for nine novae, they concluded that novae are not good standard candles. Another survey, CFHT-COVET⁵ (aimed at finding transients in the gap between novae and supernovae) found a dozen nova candidates in many galaxies in the Virgo supercluster, including some in the far outskirts of galaxies (Kasliwal et al 2010, in prep).

Here, we report on novae discovered in high cadence monitoring observations of a representative collection of galaxies with distance less than that of the Virgo cluster. The original motivation of this search — P60-FasTING⁶ was to explore rapid transients (those which last less than a couple of nights) in the nearest galaxies. A strong spectroscopic follow-up effort was a part of P60-FasTING. The survey was capable of finding novae in the major galaxies out to 4 Mpc: M31, M81, the star-burst M82 and NGC 2403. We present our sample of 21 transients, which although spectroscopically indistinguishable from classical novae, photometrically occupy a new region of

mans@astro.caltech.edu

¹ Astronomy Department, California Institute of Technology, 105-24, Pasadena, CA 91125, USA² Department of Astronomy, University of California at Berkeley, Berkeley, CA 94720, USA³ Max-Planck Institut fuer Extraterrestrische Physik, 85748 Garching, Germany⁴ We use the term candidate where the light curve is very sparse and/or there is no spectroscopic confirmation⁵ Canada France Hawaii Telescope COma Virgo Exploration for Transients⁶ Palomar 60-inch Fast Transient In Nearest Galaxies

phase space.

The paper is organized as follows: § 2 describes the discovery, photometric and spectroscopic follow-up observations of this nova sample, § 3 describes the data analysis, § 4 discusses the implications and § 5 presents our conclusion.

2. OBSERVATIONS

2.1. Experiment Design

P60-FasTING was designed with the specific goal of probing new phase space, particularly, fast transients with peak luminosity in the gap between novae and supernovae. The sample of galaxies included the brightest and nearest galaxies (<20 Mpc, majority around 10 Mpc). The survey was undertaken in a single filter (primarily Gunn-*g* and some Gunn-*i* data just around full moon). The limiting magnitude was typically Gunn-*g* <21 and cadence was <1 day. The field of view of P60 is $13.5' \times 13.5'$ and all galaxies except M31 were covered in a single pointing. For M31, five pointings were chosen to cover a larger fraction of the galaxy.

A real-time data reduction and transient search pipeline was written and implemented in April 2008. P60-FasTING ended in March 2009. The search pipeline was written in `python`. A deep reference image was constructed by combining images from several of the best seeing dark nights. Next, `wcsxmap` was used to align every new image with the reference and `hotpants` was used to compute a convolution kernel prior to image subtraction (both codes supplied by A. Becker⁷). Although the image subtraction software was quite sophisticated in its convolution of the new image to match the reference prior to subtraction, we suffered from a large number of false positives. To distill the false subtraction residuals from the bonafide astrophysical sources, a variety of automatic filters were used (e.g. the shape characteristics of the PSF of the candidate, how well does it resemble the PSF characteristics of other stars in the image). However, the final step in the vetting process was done by human eyes on candidate thumbnails every morning. Due to the myriad trade-offs for maximum completeness and minimum contamination, the complex issue of quantifying the completeness of the nova sample is beyond the scope of this publication.

Our survey was sensitive to classical novae only in a handful of the nearest galaxies in our sample (distance, $d < 4$ Mpc). Classical novae discovered by P60-FasTING are summarized in Table 1. Some novae in M31 were announced by different groups before P60-FasTING's first detection (usually due to bad weather at Palomar) — these are summarized in Table 2.

2.2. Photometry

The robotic Palomar 60-inch has a standard data reduction pipeline (Cenko et al. 2006). This pipeline performs basic detrending (flat-fielding and bias subtraction) and computes an astrometric solution. In August 2008, we added a new functionality: computation of a photometric solution. We used the SDSS catalog where available, otherwise the NOMAD catalog. Note that where NOMAD was used (e.g. M31), the transforma-

⁷ <http://www.astro.washington.edu/users/becker/c-software.html>




FIG. 1.— Location of ten novae in M31. Background image is a mosaic based on Massey data.



FIG. 2.— Location of three novae discovered by P60-FasTING in the starburst environment of M82. Background image is an HST/ACS mosaic.

tion from Johnson *UBVRI* magnitudes to SDSS *ugriz* magnitudes was done following Jordi et al. (2006).

To compute a light curve, we first measured the magnitude of the nova on the subtracted image. The subtracted image was scaled to the same flux level as the new image. Thus, we measured the magnitude of ~ 150 reference stars on the new image to compute a relative zeropoint with appropriate outlier rejection. Finally, we applied this relative zeropoint to the instrumental magnitude of the nova.

2.3. Spectroscopy

An integral part of P60-FasTING was follow-up spectroscopy to confirm and classify discovered candidate transients. Since we were looking for fast evolving phenomenon, we triggered our Target Of Opportunity pro-

TABLE 1
Novae Discovered by P60-FasTING

Nova	Host	Discovery Date	RA(J2000)	DEC(J2000)	Offset from Host	Reference
P60-NGC2403-090314	NGC2403	2009 Mar 14.160	07:36:35.00	+65:40:20.8	101.0" W, 252.0" N	Kasliwal et al. (2009a)
P60-M82OT-090314	NGC3034	2009 Mar 14.496	09:56:12.60	+69:41:32.3	104.2" E, 48.2" N	...
P60-M81OT-090213	NGC3031	2009 Feb 13.404	09:55:35.96	+69:01:51.0	15" E, 124" S	Kasliwal et al. (2009b)
P60-M31OT-081230 (2008-12b)	NGC224	2008 Dec 30.207	00:43:05.03	+41:17:52.3	233.4" E, 103.8" N	Kasliwal et al. (2009c)
P60-M81OT-081229	NGC3031	2008 Dec 29.373	09:55:38.15	+69:01:43.6	26.7" E, 131.4" S	Rau et al. (2009a)
P60-M81OT-081203	NGC3031	2008 Dec 3.303	09:55:16.92	+69:02:17.7	87.2" W, 97.4" S	Kasliwal et al. (2008a)
P60-M82OT-081119	NGC3034	2008 Nov 19.536	09:55:58.39	+69:40:56.2	29.5" E, 10.4" N	Kasliwal et al. (2008b)
P60-M81OT-081027	NGC3031	2008 Oct 27.402	09:55:36.11	+69:03:22.0	15.8" E, 33.1" S	Kasliwal et al. (2008g)
P60-M81OT-080925	NGC3031	2008 Sep 25.49	09:55:59.35	+69:05:57.1	2.35" E, 2.03" N	Kasliwal et al. (2008i)
P60-M31OT-080915 (2008-09c)	NGC224	2008 Sep 15.36	00:42:51.42	+41:01:54.0	1.34" E, 14.24" S	Kasliwal et al. (2008f)
P60-M31OT-080913 (2008-09a)	NGC224	2008 Sep 13.18	00:41:46.72	+41:07:52.1	10.8" W, 8.3" S	Kasliwal et al. (2008e)
P60-NGC891OT-080813	NGC891	2008 Aug 13.45	02:22:32.70	+42:21:56.1	8" W, 59" N	Kasliwal et al. (2008c)
P60-M31OT-080723 (2008-07b)	NGC224	2008 Jul 23.33	00:43:27.28	+41:10:03.3	8.1" E, 6.1" S	Kasliwal et al. (2008d)
P60-M82OT-080429	NGC3034	2008 Apr 29.24	09:55:21.00	+69:39:42.0	165" W, 64" S	Kasliwal et al. (2008h)
P60-M81OT-071213	NGC3031	2007 Dec 13.40	09:55:25.98	+69:04:34.8	40" W, 40" N	Kasliwal et al. (2007)

TABLE 2
Additional M31 Novae

Nova Name	Discovery Date	Classification	Reference
2007-10a	54380.606	Fe II	Pietsch et al. (2007); Gal-Yam & Quimby (2007)
2007-11f	54433.716	...	Ovcharov et al. (2007)
2007-12b	54444.528	He/N	Nakamo, Hornoch Lee et al. (2007); Bode et al. (2009)
2008-08c	54708.127	...	Valcheva et al. (2008); Hornoch
2008-10b	54759.698	Fe II	Henze et al. (2008); Di Mille et al. (2008); Barsukova et al. (2008)
2008-11a	54774.438	Hybrid	Nishiyama; Hornoch Shafter et al. (2008)



FIG. 3.— Location of one nova discovered by P60-FasTING in NGC2403. Background image is a deep co-add of P60 data.



FIG. 4.— Location of six novae discovered by P60-FasTING in M81. Background image is a color mosaic using SDSS data.

gram on the Keck I and Palomar Hale telescopes soon after discovery. Sometimes due to bad weather or bright moon-phase⁸, neither of these was an option. We resorted to the queue-scheduled service-observed programs on Gemini or HET telescopes. A log of spectroscopic observations can be found in Table 3.

We emphasize that spectroscopy is crucial in distin-

⁸ low resolution spectrographs are usually available on both these telescopes only during the dark fortnight

guishing between an optical transient which happened to be co-incident with a nearby galaxy and a classical nova. For instance, we took spectra of several optical transient candidates which did not turn out to be novae: a foreground M-dwarf flare in the Milky Way spatially coincident with NGC 7640; a background supernova; a luminous blue variable in NGC 925.




FIG. 5.— Lightcurves of novae in M31 discovered by P60-FasTING. Note the well-sampled rise.

Data were reduced in *iraf* using standard tasks in the NOAO package *onedspec* and the spectra are shown in Figure 10.

3. ANALYSIS

The primary photometric analysis was to measure the peak absolute magnitude and rate of decline. The peak magnitude had to be corrected for extinction using spectra or colors. The primary spectroscopic analysis was to classify the spectra.

3.1. Extinction

A multitude of methods have been used in the literature to measure extinction to novae. Darnley et al. (2006) compared a synthetic dust-free stellar $r - i$ map of M31 to an observed $r - i$ color map of M31 and used the difference between the maps to generate a dust map of M31. The location of the nova on this map determined how much extinction needed to be applied. This assumed that the novae were behind the galaxy and suffered extinction due to the entire column of dust. The average extinction as determined by this method is $A_i=0.8$. The galactic extinction along the line of sight of M31 of $A_i=0.13$. Shafter et al. (2009) compared the observed color of the new nova to that of a well-studied nova to derive the extinction. Kogure (1961) used the Balmer

decrement (H_α/H_β) and attributed the excess in the ratio over the theoretical Case B value to dust.

Our preferred method of computing extinction is by using the spectroscopic Balmer decrement where nebular spectra are available. We subtract the continuum, measure the flux ratio of the two lines, and then use:

$$\begin{aligned} A_g &= 3.793 \times E(B - V) = 3.650 \times E(g - r) \\ &= 3.650 \times \log_{10}(H_\alpha/H_\beta) - 1.75 \pm 0.14 \end{aligned}$$

The uncertainty comes from the range in expected ratios for Case B of 2.76–3.30.

Our second choice is to use the $g - i$ color of the nova at maximum, compare against the typical $g - i$ color and attribute the reddening to dust.

van den Bergh & Younger (1987) compiled photometry of several Galactic novae and derived an average color at maximum of $\langle B - V \rangle_0 = 0.23 \pm 0.06$ mag. Following Shafter et al. (2009), we can use the colors of an A5V star ($T=8200$ K) to translate $\langle B - V \rangle_0$ to $\langle g - i \rangle_0$. Using colors of an A5V star from Kraus & Hillenbrand (2007), we get:

$$\langle g - i \rangle_0 = 1.88 - 2.15 = -0.27$$

mag. Now,

$$A_g = 3.793 \times E(B - V) = 2.223 \times E(g - i)$$




FIG. 6.— Lightcurves of novae in M81.

. Hence,

$$A_g = 2.223 \times [(g-i)_{obs} - \langle g-i \rangle_0] = 2.223 \times [(g-i)_{obs} + 0.27]$$

For the case of M31N2007-11d (Shafter et al. 2009), where data for both the above options are available, we get consistent answers: $g-i=0.7$ mag at maximum suggests $A_g=0.75$ mag and a Balmer decrement of 4.6 suggests $A_g=0.65$ mag.

If neither a color at maximum nor a spectrum at late-

time is available, we use the average line-of-sight extinction to the host galaxy using the Schlegel maps. The uncertainty in extinction calculation significantly contributes to the uncertainty in the peak magnitude.

We note here that for the case of P60-M82OT-081119, the light curve was unusually red for a nova and the extinction correction may be overestimated.

3.2. Rate of decline




FIG. 7.— Lightcurves of novae in M82 and NGC2403. Note that P60-M82OT-081119 is much redder than typical novae.

The heterogeneity in nova light curves suggests that a single parameter may not characterize the decline well. Traditionally, the time to decay from peak by one magnitude (t_1), two magnitudes (t_2) or three magnitudes (t_3) is used. For several novae (e.g. M31N2007-10a, M31N2008-08c, M31N2008-11a), the decline is more or less linear and t_1 can be approximated as half of t_2 . For some novae (e.g. M31N2008-10b), the light curve behavior is more complex and this simplification is not applicable. In Table 4, we see that t_2 values (where available) are sometimes larger and sometimes smaller than twice the t_1 value.

We note here that we did not have data to measure the decline of the nova in NGC 891 and hence it is excluded from further MMRD analysis.

3.3. Rate of rise

Given the cadence of P60-FastTING, we were able to catch several novae on the rise. We define the rate of rise as the average slope between first detection and peak detection and summarize in Table 4. We find a wide range of rise-times, from $>1.8 \text{ mag day}^{-1}$ (e.g. M31N2008-11a) to 0.2 mag day^{-1} (e.g. M31N2008-09a). It is not clear how previous determinations of the MMRD in the literature dealt with the uncertainty in the peak magnitude due to inadequate coverage. Especially since previous

surveys likely had a relatively slower cadence, missing the peak may be a substantial source of error. Slower cadence and/or shallower depth would correspond to a weaker constraint on the rise time of the nova. Due to gaps on account of weather, some of the P60 light curves have constraint weaker than 0.1 mag day^{-1} on the rate of rise. Hence, we do not use the lightcurves of P60-M81OT-080926 or P60-M82OT-080429 for subsequent analysis of the MMRD relation.

3.4. Spectral Classification

For spectroscopy the primary analysis was to classify the novae by their spectra. The taxonomy of novae were laid out by Payne-Gaposchkin (1964) and McLaughlin (1960). The most prominent feature in all classical novae is Balmer emission. Williams (1992) propose that there is a two component structure of the emitting gas – discrete shell and continuous wind. If the wind mass-loss rate is low, the effective photosphere is smaller, the radiation temperature higher, the level of ionization of the shell higher, resulting in a shell-dominated He/N spectrum. If the wind mass-loss rate is high, it results in a wind-dominated Fe II spectrum.

Thus, classical novae are divided into two principal families — the “Fe” class (dominated by Fe II lines, often




FIG. 8.— Lightcurves of additional novae in M31.

low velocity and showing P-Cygni profiles) and “He/N” class (dominated by He and N lines, often high velocity and flat or jagged-topped profiles). These evolve into nebular spectra with four classes based on the prominent forbidden lines — standard (e.g. [N II], [O II], [O III]), neon (e.g. [Ne V], [Ne III]), coronal (e.g. [Fe X]) or no forbidden lines. The Fe class novae are expected to evolve into standard or neon nebular spectra. The He/N class are expected to evolve into neon, coronal or

no forbidden line spectra. Some novae are classified as “hybrid” as they start out with high velocity Fe II features and quickly evolve into showing He/N features (e.g. V745 Sco, V3890 Sgr, M31N2008-11a).

Majority of the P60-FasTING spectra show clear permitted lines from Fe II (42), Fe II (37,38) and O I. The line velocities are low and typical Gaussian FWHM are $< 2500 \text{ km s}^{-1}$ with the exception of P60-M81-080925 where H_{α} velocity is 3000 km s^{-1} . P60-NGC2403-090314

TABLE 3
Spectroscopy

Transient Name	Spectroscopy Date	Telescope	Instrument	Classification	Observer
P60-NGC2403-090314	2009 Mar 20.145	P200	DBSP (Oke & Gunn 1982)	Fe Class?	Kasliwal, Ellis
P60-M81-090213	2009 Feb 18.510	Keck I	LRIS (Oke et al. 1995)	Fe Class	Ofek
P60-M81-081229	2008 Dec 31.40	P200	DBSP (Oke & Gunn 1982)	Fe Class	Rau, Salvato
P60-M31-081230	2008 Dec 31.104	P200	DBSP (Oke & Gunn 1982)	Fe Class	Rau, Salvato
P60-M81-081203	2008 Dec, 4, 5, 16	P200, Gemini	DBSP, GMOS-N (Hook et al. 2004)	Fe Class	Kasliwal
P60-M81-080925	2008 Sep 29.51	P200	DBSP (Oke & Gunn 1982)	Fe Class	Quimby
P60-M31-080915	2008 Sep 20.2	HET	LRS (Hill et al. 1998)	Fe Class	Shafter
P60-M31-080913	2008 Sep 22.4	HET	LRS (Hill et al. 1998)	Fe Class	Shafter
P60-M31-080723	2008 Aug 1	P200	DBSP (Oke & Gunn 1982)	Fe Class	Ofek
P60-M82-080429	2008 May 2.28	P200	DBSP (Oke & Gunn 1982)	...	Cenko
P60-M81-071213	2007 Dec 15.565	Keck	LRIS (Oke et al. 1995)	...	Ofek

TABLE 4
Nova Characteristics

Nova Name	Balmer Decrement $FH\alpha/H\beta$	Spectral Phase	Color at Peak $g - i(\text{mag})$	Extinction $A_g(\text{mag})$	Rate of Rise mag day^{-1}	Abs-Mag $M_g(\text{mag})$	t1 days	t2 days
P60-NGC2403-090314	5.0	Nebular	...	0.8	1.3	-9.0	3.3	>6
P60-M82-090314	...	...	-0.3	0.6	>1.2	-8.5	2.4?	>3.3
P60-M81-090213	...	Nebular	0.7	2.2	0.2	-9.9	5?	>10.9
P60-M31-081230 (2008-12b)	...	...	-0.17	0.24	0.6	-7.5	12.3	
P60-M81-081229	2.4	Near-Max	0.1	0.90	>0.1	-8.7	2.9	...
P60-M81-081203	...	...	-0.03	0.53	0.7	-8.0	7.5?	>23
P60-M82-081119	...	...	0.98	2.8	0.6	-10.7	4.0	...
P60-M81-081027	...	...	...	0.3	0.2	-7.6	4.0	...
P60-M81-080926	1.4	Near-Max	-0.2	0.13	...	-8.5	8.9	14.0
P60-M31-080915 (2008-09c)	1.4	Near-Max	-0.72	0.24	0.4	-7.8	9.1	16.6
P60-M31-080913 (2008-09a)	2.5	Near-Max	-0.30	0.24	0.2	-6.8	6.3	16.0
P60-M31-080723 (2008-07b)	14.3	Nebular?	-0.20	2.5	0.2	-7.6	5.0	12.0
P60-M82-080429	6.3	Nebular?	...	1.2	...	-8.5	8.1	...
P60-M81OT-071213	3.8	Nebular?	0.5	0.4	>0.6	-7.8	1.0	...
2007-10a	...	...	...	>0.24	>1.2	-7.0	4.1?	8.6
2007-11f	...	...	-0.16	0.24	>0.1	-5.1	5.0	>8.0
2007-12b	...	...	...	>0.24	>0.6	-6.3	3.5	>5.0
2008-08c	...	...	...	>0.24	0.3	-7.5	11.0	26.3
2008-10b	...	...	-0.58	>0.24	0.2	-6.5	6.0	12.3?
2008-11a	...	...	0.41	1.5	>1.8	-7.7	2.9	7.5

shows weak Fe II(42) and weak P-Cygni profiles in Balmer lines and can tentatively also be classified as Fe II class. P60-M81-071213 and P60-M82-080429 have very low SNR and no feature other than the Balmer lines are detected, hence, we cannot classify them. Multiple spectra of P60-M81-081203 were taken – initially, the spectra show a featureless continuum (obtained a few days prior to maximum light) and later (about a week after maximum light), evolved to show Balmer lines, Fe II (42), O I. We summarize spectral classifications in Table 3. For four novae in M31, other groups obtained spectra and we summarize their classification in Table 2.

4. DISCUSSION

In comparison to traditional nova searches, P60-FastTING was designed as a faster cadence and deeper survey. Weather-permitting, galaxies in the sample were imaged every night to a mean depth of $G_{\text{unn}} < 21$ mag. Hence, P60-FastTING was sensitive to tran-

sients that are less luminous and evolve faster than classical novae.

Given that our light curves are well-sampled, and we have spectra or color measurements to correct for extinction, we can securely measure both the maximum magnitude and the rate of decline. To our surprise, as demonstrated in Figure 11, we find that the P60-FastTING nova sample is evidently inconsistent with the MMRD relation (della Valle & Livio 1995).

In Figure 11, the decay time is measured as the time to decay by one magnitude. To test whether the apparent photometric diversity is consistent with the MMRD over a longer timescale, we plot the time to decay by two magnitudes in Figure 12. Furthermore, we restrict this to the sub-sample of six classical novae in M31 with the best-sampled light curves (see lightcurves of P60-M31OT-080915, P60-M31OT-080913, P60-M31OT-080723, M31N2007-10a, M31N2008-08c, M31N2008-11a in Figure 5 and Figure 8). Even this sub-sample does




FIG. 9.— Lightcurve of possible nova in NGC 891. This was not spectroscopically confirmed and is not used in the MMRD analysis.

not obey the MMRD relation. This scatter is larger than the ± 0.8 mag predicted on theoretical grounds by Shara (1981).

Despite the atypical photometric signature, the P60-FasTING nova sample shows no spectroscopic peculiarities. In Figure 11, the symbols indicate the spectral class — majority are Fe II class (stars), a couple are He/N class (circles), some have spectra with no prominent features for classification (filled squares) and a few have no spectra (empty squares).

We could hypothesize that some of the P60-FasTING novae are not classical but recurrent (classical novae which recur on a timescales shorter than a century) since recurrent novae are also known not to obey the MMRD relation. Recurrent novae are expected to occur in the most massive white dwarfs with high accretion rates. A small amount of mass accreted on a short timescale is sufficient to trigger thermonuclear runaway. Recently, Schaefer (2009) compiled all available photometry over the past century on the ten recurrent novae in our galaxy — overplotted to compare with the P60-FasTING sample in Figure 13.

There are three ways to test the recurrent nova hypothesis. First, spectra of recurrent novae have high velocities and belong to either He/N or hybrid class (e.g. V3890 Sgr, V745 Sco, V394 CrA in Williams 1992 and Williams et al. 1994). We find that majority off the P60-FasTING novae do not share the spectroscopic properties of recurrent novae. P60-FasTING novae mostly belong to the Fe II class and have low velocities. Second, recurrent novae are often a few magnitudes brighter than classical novae at quiescence. Schaefer (2009) suggest that recurrent novae range from $-4.1 < M_V < 3.2$ and classical novae range from $1.1 < M_V < 7.0$ at quiescence. Given the distance modulus to these galaxies, this test is within reach of 10-m class telescopes and easy with HST (e.g. see Bode et al. 2009). Third, the unambiguous test of whether an eruption is recurrent is to continue to monitor these galaxies for the next few decades until another eruption is witnessed.

In order to decipher the nature of this new sub-class of novae we turn to the fundamental physics of classical novae. The physics is governed by four parameters

- mass of the white dwarf, temperature, accretion rate and composition. The MMRD relation is explained with the mass of the white dwarf being the single, dominant parameter. Perhaps, the P60-FasTING sample of faint and fast novae can be explained based on an unexplored region of this four-parameter phase space? Could some P60-FasTING novae come from hot and massive white dwarfs? If it is hot, then the thermonuclear runaway would not be as explosive and thus, the peak luminosity would be fainter. Also, the higher temperature would result in a smaller amount of envelope mass being sufficient to trigger thermonuclear runaway and thus, the timescale would be faster.

Recent theoretical efforts have explored nova diversity (e.g. Townsley & Bildsten 2004, Shen & Bildsten 2009, Epelstain et al. 2007). Yaron et al. (2005) present an extended grid of nova models to explore a wider parameter space (in mass, temperature and accretion rate) than traditionally explored for classical novae subject to physical constraints (such as conditions for thermonuclear runaway). We summarize the results of the variety of models they run in Figure 14. Some hot and massive white dwarfs with high accretion rates can result in a faint and fast nova population consistent with the P60-FasTING sample. Indeed, Yaron et al. (2005) predict the existence of remarkably small amplitude novae across the entire span of decay rates.

Finally, we note that more than half of the P60-FasTING nova sample is inconsistent with the MMRD. This suggests that faint and fast novae are commonplace and cannot be explained by a rare type of white dwarf.

5. CONCLUSION

We conclude that P60-FasTING has uncovered classical novae in a new region in the luminosity-timescale phase space of optical transients. Classical novae span at least two orders of magnitude in time and two orders of magnitude in luminosity. Future surveys would have a large enough sample to meaningfully constrain the relative populations of classical novae in the different areas of phase space.

P60-FasTING was designed as a pilot project, to begin to set the stage for future projects such as Palomar Transient Factory (PTF⁹, Law et al. 2009, Rau et al. 2009b, Rahmer et al. 2008), PanSTARRS (PS1¹⁰) and Large Synoptic Survey Telescope (LSST¹¹). Both PTF and PS1 are now underway. PTF is looking at several nearby galaxies with a similar depth and cadence as P60-FasTING. Among nearby galaxies, PS1 day-cadence fields only cover M31 but are a couple of magnitudes deeper. LSST will be both deeper and faster cadence and cover the visible sky. P60-FasTING is only the trailblazer for the uncovering of a wealth of information about classical novae by near-future synoptic surveys.

We thank Marina Orio for discussion, in particular, the suggestion that some of the faint novae may be recurrent novae. We thank A. Shafter, M. Shara, L. Bild-

⁹ <http://www.astro.caltech.edu/ptf>

¹⁰ <http://pan-starrs.ifa.hawaii.edu>

¹¹ <http://www.lsst.org>




FIG. 10.— Optical spectra of P60-FasTING novae. Majority are Fe II class. Note that they have been arbitrarily offset along the y-axis for clarity.




FIG. 11.— Maximum magnitude (absolute g -band) versus rate of decline (time to decay from peak by 1 mag). Gray region denotes the della Valle & Livio (1995) MMRD relation and dark gray dots the nova sample they used. The P60-FasTING sample is shown with symbols that denote spectral type — Fe II class (star), He/N class (circle), spectrum with no prominent features for classification (squares) and no spectrum (empty square).




FIG. 12.— Maximum magnitude (absolute g -band) versus rate of decline (time to decay from peak by 2 mag). Only the six novae with the best sampled light curves are shown.

sten and O. Yaron for valuable feedback. We are grateful to A. Becker for making his software `hotpants` and `wcsremap` available for public use. MMK thanks the Gordon and Betty Moore Foundation for the Hale Fellowship in support of graduate study. S.B.C. is grateful for generous support from Gary and Cynthia Bengier, the Richard and Rhoda Goldman Fund, and National Science Foundation (NSF) grant AST0908886. MMK thanks the Palomar Observatory staff for their help in maximizing the efficiency and image quality of the Palomar 60-inch. Some of the data presented herein were obtained at the W.M. Keck Observatory, which is operated




FIG. 13.— Comparison of the P60-FasTING nova sample (green symbols) with the Galactic recurrent novae (blue circles, data from Schaefer (2009)).




FIG. 14.— Theoretical results of Yaron et al. (2005) over a wide range of nova parameters — we use the colors of an A5V star to convert $L_{4\text{max}}$ to M_g and approximate t_1 as $t_{\text{ml}}/3$. The size of the symbol is proportional to the mass of the white dwarf. The color of the symbol denotes temperature — 10 million K (red), 30 million K (green), 50 million K (blue). Empty circles denote lower accretion rate in the range $10^{-12.3}$ – $10^{-10} M_{\odot} \text{yr}^{-1}$ and filled circles denote higher accretion rate in the range 10^{-9} – $10^{-6} M_{\odot} \text{yr}^{-1}$. Note that the density of circles is unrelated to the relative populations.

as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation. The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity

to conduct observations from this mountain.

REFERENCES

- Arp, H. C. 1956, *AJ*, 61, 15
- Barsukova, E., Fabrika, S., Hornoch, K., Sholukhova, O., & Valeev, A. 2008, *The Astronomer's Telegram*, 1871, 1
- Bode, M. F., Darnley, M. J., Shafter, A. W., Page, K. L., Smirnova, O., Anupama, G. C., & Hilton, T. 2009, *ApJ*, 705, 1056
- Darnley, M. J., Bode, M. F., Kerins, E., Newsam, A. M., An, J., Baillon, P., Belokurov, V., Calchi Novati, S., Carr, B. J., Cr    , M., Evans, N. W., Giraud-H  raud, Y., Gould, A., Hewett, P., Jetzer, P., Kaplan, J., Paulin-Henriksson, S., Smartt, S. J., Tsapras, Y., & Weston, M. 2006, *MNRAS*, 369, 257
- della Valle, M. & Livio, M. 1995, *ApJ*, 452, 704
- Di Mille, F., Ciroi, S., Orio, M., Rafanelli, P., Bianchini, A., Nelson, T., & Andreuzzi, G. 2008, *The Astronomer's Telegram*, 1818, 1
- Downes, R. A. & Duerbeck, H. W. 2000, *AJ*, 120, 2007
- Epelstain, N., Yaron, O., Kovetz, A., & Prialnik, D. 2007, *MNRAS*, 374, 1449
- Ferrarese, L., C    , P., & Jord  n, A. 2003, *ApJ*, 599, 1302
- Gal-Yam, A. & Quimby, R. 2007, *The Astronomer's Telegram*, 1236, 1
- Henze, M., Pietsch, W., Burwitz, V., Hatzidimitriou, D., Reig, P., Primak, N., & Papamastorakis, G. 2008, *The Astronomer's Telegram*, 1790, 1
- Hill, G. J., Nicklas, H. E., MacQueen, P. J., Mitsch, W., Wellem, W., Altmann, W., Wesley, G. L., & Ray, F. B. 1998, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 3355, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, ed. S. D'Odorico, 433–443
- Hook, I. M., J  rgensen, I., Allington-Smith, J. R., Davies, R. L., Metcalfe, N., Murowinski, R. G., & Crampton, D. 2004, *PASP*, 116, 425
- Hornoch, K., Scheirich, P., Garnavich, P. M., Hameed, S., & Thilker, D. A. 2008, *A&A*, 492, 301
- Hubble, E. P. 1929, *ApJ*, 69, 103
- Jordi, K., Grebel, E. K., & Ammon, K. 2006, *A&A*, 460, 339
- Kasliwal, M. M., Cenko, S. B., Ofek, E. O., Quimby, R., Rau, A., & Caltech, S. R., K. 2009a, *The Astronomer's Telegram*, 1984, 1
- Kasliwal, M. M., Cenko, S. B., Ofek, E. O., Quimby, R., Rau, A., & Kulkarni, S. R. 2009b, *The Astronomer's Telegram*, 1934, 1
- Kasliwal, M. M., Cenko, S. B., Quimby, R., Rau, A., Ofek, E. O., & Kulkarni, S. R. 2008a, *The Astronomer's Telegram*, 1880, 1
- Kasliwal, M. M., Cenko, S. B., Rau, A., Ofek, E. O., Quimby, R., & Kulkarni, S. R. 2008b, *The Astronomer's Telegram*, 1854, 1
- , 2008c, *The Astronomer's Telegram*, 1663, 1
- , 2008d, *The Astronomer's Telegram*, 1627, 1
- , 2008e, *The Astronomer's Telegram*, 1717, 1
- , 2008f, *The Astronomer's Telegram*, 1719, 1
- , 2008g, *The Astronomer's Telegram*, 1826, 1
- Kasliwal, M. M., Ofek, E. O., Rau, A., Cenko, S. B., Quimby, R., & Kulkarni, S. R. 2008h, *The Astronomer's Telegram*, 1646, 1
- Kasliwal, M. M., Quimby, R., Cenko, S. B., Rau, A., Ofek, E. O., & Kulkarni, S. R. 2008i, *The Astronomer's Telegram*, 1764, 1
- Kasliwal, M. M., Quimby, R., Rau, A., Ofek, E., Cenko, B., & Kulkarni, S. 2007, *The Astronomer's Telegram*, 1330, 1
- Kasliwal, M. M., Rau, A., Salvato, M., Cenko, S. B., Ofek, E. O., Quimby, R., & Kulkarni, S. R. 2009c, *The Astronomer's Telegram*, 1886, 1
- Kogure, T. 1961, *PASJ*, 13, 335
- Kraus, A. L. & Hillenbrand, L. A. 2007, *AJ*, 134, 2340
- Law, N. M., Kulkarni, S. R., Dekany, R. G., Ofek, E. O., Quimby, R. M., Nugent, P. E., Surace, J., Grillmair, C. C., Bloom, J. S., Kasliwal, M. M., Bildsten, L., Brown, T., Cenko, S. B., Ciardi, D., Cr  ner, E., Djorgovski, S. G., van Eyken, J., Filippenko, A. V., Fox, D. B., Gal-Yam, A., Hale, D., Hamam, N., Helou, G., Henning, J., Howell, D. A., Jacobsen, J., Laher, R., Mattingly, S., McKenna, D., Pickles, A., Poznanski, D., Rahmer, G., Rau, A., Rosing, W., Shara, M., Smith, R., Starr, D., Sullivan, M., Velur, V., Walters, R., & Zolkower, J. 2009, *PASP*, 121, 1395
- Lee, C., Ries, C., Riffeser, A., & Seitz, S. 2007, *The Astronomer's Telegram*, 1324, 1
- Livio, M. 1992, *ApJ*, 393, 516
- McLaughlin, D. B. 1960, in *Stellar Atmospheres*, ed. J. L. Greenstein, 585–
- Oke, J. B., Cohen, J. G., Carr, M., Cromer, J., Dingizian, A., Harris, F. H., Labrecque, S., Lucinio, R., Schaaf, W., Epps, H., & Miller, J. 1995, *PASP*, 107, 375
- Oke, J. B. & Gunn, J. E. 1982, *PASP*, 94, 586
- Ovcharov, E., Valcheva, A., Kostov, A., Nikolov, Y., Georgiev, T., & Nedialkov, P. 2007, *The Astronomer's Telegram*, 1312, 1
- Payne-Gaposchkin, C. 1964, *The galactic novae*, ed. Gaposchkin, C. H. P.
- Pietsch, W., Burwitz, V., Stoss, R., Updike, A., Hartmann, D., Milne, P., & Williams, G. 2007, *The Astronomer's Telegram*, 1230, 1
- Rahmer, G., Smith, R., Velur, V., Hale, D., Law, N., Bui, K., Petrie, H., & Dekany, R. 2008, in *Presented at the Society of Photo-Optical Instrumentation Engineers (SPIE) Conference*, Vol. 7014, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*
- Rau, A., Kasliwal, M. M., & Salvato, M. 2009a, *The Astronomer's Telegram*, 1887, 1
- Rau, A., Kulkarni, S. R., Law, N. M., Bloom, J. S., Ciardi, D., Djorgovski, G. S., Fox, D. B., Gal-Yam, A., Grillmair, C. C., Kasliwal, M. M., Nugent, P. E., Ofek, E. O., Quimby, R. M., Reach, W. T., Shara, M., Bildsten, L., Cenko, S. B., Drake, A. J., Filippenko, A. V., Helfand, D. J., Helou, G., Howell, D. A., Poznanski, D., & Sullivan, M. 2009b, *PASP*, 121, 1334
- Schaefer, B. E. 2009, *ArXiv e-prints*
- Shafter, A. W., Ciardullo, R., Bode, M. F., Darnley, M. J., & Misselt, K. A. 2008, *The Astronomer's Telegram*, 1832, 1
- Shafter, A. W., Rau, A., Quimby, R. M., Kasliwal, M. M., Bode, M. F., Darnley, M. J., & Misselt, K. A. 2009, *ApJ*, 690, 1148
- Shara, M. M. 1981, *ApJ*, 243, 926
- Shen, K. J. & Bildsten, L. 2009, *ApJ*, 692, 324
- Townsend, D. M. & Bildsten, L. 2004, *ApJ*, 600, 390
- Valcheva, A., Ovcharov, E., Latev, G., Kostov, A., Nikolov, Y., Georgiev, T., & Nedialkov, P. 2008, *The Astronomer's Telegram*, 1687, 1
- van den Bergh, S. & Younger, P. F. 1987, *A&AS*, 70, 125
- Williams, R. E. 1992, *AJ*, 104, 725
- Williams, R. E., Phillips, M. M., & Hamuy, M. 1994, *ApJS*, 90, 297
- Yaron, O., Prialnik, D., Shara, M. M., & Kovetz, A. 2005, *ApJ*, 623, 398
- Zwicky, F. 1936, *PASP*, 48, 191